La Primera Medida de la Sección Eficaz Doble Diferencial en la Dispersión Quasi-elastica de la Corriente Cargada del Neutrino Muónico

Work based on PhD thesis at Indiana University

Teppei Katori for the MiniBooNE collaboration
Massachusetts Institute of Technology
NuInt 09, Sitges, May, 19, 09

Massachusetts Institute of Technology

First Measurement of Muon Neutrino Charged Current Quasielastic (CCQE) Double Differential Cross Section

outline

1. Booster neutrino beamline
2. CCQE events in MiniBooNE
3. CC1π background constraint
4. CCQE M_A^{eff}-κ shape-only fit
5. CCQE absolute cross section
6. Conclusion

Work based on PhD thesis at Indiana University

Teppei Katori for the MiniBooNE collaboration
Massachusetts Institute of Technology
NuInt 09, Sitges, May, 19, 09

Massachusetts Institute of Technology

1. Booster neutrino beamline

2. CCQE events in MiniBooNE

3. CC1π background constraint

4. CCQE M_A^{eff}-κ shape-only fit

5. CCQE absolute cross section

6. Conclusion

1. Booster Neutrino Beamline



MiniBooNE extracts 8.9 GeV/c momentum proton beam from the Booster



1. Booster Neutrino Beamline

Magnetic focusing horn



Protons are delivered to a beryllium target in a magnetic horn (flux increase ~6 times)



1. Booster Neutrino Beamline

Modeling of meson production is based on the measurement done by HARP collaboration

- Identical, but 5% λ Beryllium target
- 8.9 GeV/c proton beam momentum

HARP collaboration, Eur.Phys.J.C52(2007)29

HARP experiment (CERN)



Booster neutrino beamline pion kinematic space



Majority of pions create neutrinos in MiniBooNE are directly measured by HARP (>80%)

1. Booster Neutrino Beamline

Modeling of meson production is based on the measurement done by HARP collaboration

- Identical, but 5% λ Beryllium target
- 8.9 GeV/c proton beam momentum

HARP collaboration, Eur.Phys.J.C52(2007)29

HARP experiment (CERN)



HARP data with 8.9 GeV/c proton beam momentum



The error on the HARP data (~7%) directly propagates.
The neutrino flux error is the dominant source of normalization error for an absolute cross section in MiniBooNE.

1. Booster Neutrino Beamline



The decay of mesons make the neutrino beam. The neutrino beam is dominated by ν_μ (93.6%), of this, 96.7% is made by π^+-decay

$$\pi^+ \rightarrow \mu^+ + \nu_\mu$$

MiniBooNE collaboration, PRD79(2009)072002



1. Booster neutrino beamline

2. CCQE events in MiniBooNE

3. CC1π background constraint

4. CCQE M_A^{eff}-κ shape-only fit

5. CCQE absolute cross section

6. Conclusion

2. CCQE event measurement in MiniBooNE

ν_μ charged current quasi-elastic (ν_μ CCQE) interaction is an important channel for the neutrino oscillation physics and the most abundant (~40%) interaction type in MiniBooNE detector

$$\nu_\mu + n \rightarrow p + \mu^-$$

$$(\nu_\mu + {}^{12}C \rightarrow X + \mu^-)$$



MiniBooNE detector

(spherical Cherenkov detector)

MiniBooNE collaboration, NIM.A599(2009)28



2. CCQE event measurement in MiniBooNE

ν_μ charged current quasi-elastic (ν_μ CCQE) interaction is an important channel for the neutrino oscillation physics and the most abundant (~40%) interaction type in MiniBooNE detector

$$\nu_\mu + n \rightarrow p + \mu^-$$

$$(\nu_\mu + {}^{12}C \rightarrow X + \mu^-)$$



MiniBooNE detector

(spherical Cherenkov detector)

MiniBooNE collaboration, NIM.A599(2009)28



muon like Cherenkov light and subsequent decayed electron (Michel electron) like Cherenkov light are the signal of CCQE event

proton measurement in neutral current elastic, see D. Perevalov and R. Tayloe's talk, May 20 (Wed.)

2. CCQE event measurement in MiniBooNE

ν_μ CCQE interactions (ν+n → μ+p) has characteristic two "subevent" structure from muon decay

$$\nu_\mu + n \rightarrow \overset{1}{\boxed{\mu^-}} + p \rightarrow \nu_\mu + \overline{\nu_e} + \overset{2}{\boxed{e^-}} + p$$



26.5% efficiency
75.8% purity
146,070 events
with 5.58E20POT

2. CCQE event measurement in MiniBooNE

All kinematics are specified from 2 observables, muon energy E_μ and muon scattering angle θ_μ

Energy of the neutrino E_ν^{QE} and 4-momentum transfer Q^2_{QE} can be reconstructed by these 2 observables, under the assumption of CCQE interaction with bound neutron at rest ("QE assumption")

$$E_\nu^{QE} = \frac{2(M - E_B)E_\mu - (E_B^2 - 2ME_B + m_\mu^2 + \Delta M^2)}{2[(M - E_B) - E_\mu + p_\mu \cos\theta_\mu]}$$

$$Q^2_{QE} = -m_\mu^2 + 2E_\nu^{QE}(E_\mu - p_\mu \cos\theta_\mu)$$



1. Booster neutrino beamline

2. CCQE events in MiniBooNE

3. CC1π background constraint

4. CCQE M_A-κ shape-only fit

5. CCQE absolute cross section

6. Conclusion

3. CC1π background constraint, introduction

data-MC comparison, in 2 subevent sample (absolute scale)

Problem 1

CCQE sample shows good agreement in shape, because we tuned relativistic Fermi gas (RFG) parameters.

MiniBooNE collaboration, PRL100(2008)032301

However absolute normalization does not agree.

The background is dominated with CC1π without pion (CCQE-like). We need a background prediction with an absolute scale.

CCQE

$$\nu_\mu + n \rightarrow \overset{1}{\boxed{\mu^-}} + p \rightarrow \nu_\mu + \overline{\nu_e} + \overset{2}{\boxed{e^-}} + p$$

CC1π

$$\nu_\mu + N \rightarrow \overset{1}{\boxed{\mu^-}} + \cancel{\pi^+} + N \rightarrow \nu_\mu + \overline{\nu_e} + \overset{2}{\boxed{e^-}} + N$$

(π-absorption)



3. CC1π background constraint, introduction

data-MC comparison, in 3 subevent sample (absolute scale)

Problem 2

CC1π sample is worse situation, data and MC do not agree in shape nor normalization.

Under this situation, we cannot use CC1π prediction for background subtraction for CCQE absolute cross section measurement.



CC1π

$$\nu_\mu + N \rightarrow \overset{1}{\boxed{\mu^-}} + \pi^+ + N \rightarrow \nu_\mu + \overline{\nu}_e + \overset{2}{\boxed{e^-}} + N$$

$$\hookrightarrow \nu_\mu + \mu^+ \rightarrow \nu_\mu + \overline{\nu}_e + \overset{3}{\boxed{e^+}} + N$$

recent development of prediction in CC1π, see J. Novak's talk, May 22 (Fri.),

pion measurement in CC1π, see M. Wilking's talk, May 22 (Fri.),

3. CC1π background constraint

data-MC comparison, before CC1π constraint (absolute scale)

Solution

Use data-MC Q^2 ratio in CC1π sample to correct all CC1π events in MC.

Then, this "new" MC is used to predicts CC1π background in CCQE sample

This correction gives both CC1π background normalization and shape in CCQE sample



3. CC1π background constraint

data-MC comparison, after CC1π constraint (absolute scale)

Now we have an absolute prediction of CC1π background in CCQE sample.

We are ready to measure the absolute CCQE cross section!



1. Booster neutrino beamline

2. CCQE events in MiniBooNE

3. CC1π background constraint

4. CCQE M_A^{eff}-κ shape-only fit

5. CCQE absolute cross section

6. Conclusion

4. Pauli blocking parameter “kappa”, κ

We performed shape-only fit for Q^2 distribution to fix CCQE shape within RFG model, by tuning M_A^{eff} (effective axial mass) and κ

Pauli blocking parameter "kappa”, κ

Smith and Moniz, Nucl.,Phys.,B43(1972)605

To enhance the Pauli blocking at low Q^2, we introduced a new parameter κ, which is the energy scale factor of lower bound of nucleon sea in RFG model in Smith-Moniz formalism, and controls the size of nucleon phase space

$$\mathrm{Elo} = \kappa\left(\sqrt{(p_F^2 + M^2)} - w + E_B\right)$$



4. M_A^{eff}-κ shape-only fit

M_A^{eff} - κ shape-only fit result

M_A^{eff} = 1.35 ± 0.17 GeV (stat+sys)
κ = $1.007^{+0.007}_{-\infty}$ (stat+sys)
χ^2/ndf = 47.0/38

M_A^{eff} goes even up, this is related to our new background subtraction.

κ goes down due to the shape change of the background. Now κ is consistent with 1.
κ doesn't affects cross section below ~0.995.

M_A^{eff} only fit (M_A^{eff} = 1.37 ± 0.12 GeV, χ^2/ndf = 48.6/39)

data-MC Q^2 comparison before and after fit



Fit parameter space



4. M_A^{eff}-κ shape-only fit

MiniBooNE anti-neutrino CCQE data
J. Grange poster, May 19 (Tue.)

M_A^{eff} - κ shape-only fit result

M_A^{eff} = 1.35 ± 0.17 GeV (stat+sys)
$\kappa = 1.007^{+0.007}_{-\infty}$ (stat+sys)

This new CCQE model doesn't affect our cross section result.

Data-MC agreement in T_μ-cosθ kinematic plane is good.

World averaged RFG model
M_A^{eff} = 1.03, κ = 1.000



data-MC ratio in T_μ-cosθ kinematic plane after fit



1. Booster neutrino beamline

2. CCQE events in MiniBooNE

3. CC1π background constraint

4. CCQE M_A-κ shape-only fit

5. CCQE absolute cross section

6. Conclusion

5. CCQE absolute cross section

Flux-averaged single differential cross section (Q^2_{QE})

The data is compared with various RFG model with neutrino flux averaged.

Compared to the world averaged CCQE model (red), our CCQE data is 35% high

Our model extracted from shape-only fit has better agreement (within our total normalization error).



5. CCQE absolute cross section

Flux-unfolded total cross section (E_ν^{RFG})

New CCQE model is tuned from shape-only fit in Q^2, and it also describes total cross section well.



5. CCQE errors

Error summary (systematic error dominant)

Flux error dominates the total normalization error.

Cross section error is small because of high purity and in situ background measurement.

Detector error dominates shape error, because this is related with energy scale.

Unfolding error is the systematic error associated to unfolding.



5. QE cross section comparison with NOMAD

Flux-unfolded total cross section (E_ν^{RFG})

New CCQE model is tuned from shape-only fit in Q^2, and it also describes total cross section well.

Comparing with NOMAD, MiniBooNE cross section is 35% higher, but these 2 experiments leave a gap in energy to allow some interesting physics.



5. CCQE total cross section model dependence

Flux-unfolded total cross section (E_ν^{RFG})

Unfortunately, flux unfolded cross section is model dependent.

Reconstruction bias due to QE assumption is corrected under "RFG" model assumption.

One should be careful when comparing flux-unfolded data from different experiments.



5. CCQE total cross section model dependence

Flux-unfolded total cross section (E_ν^{RFG})

Unfortunately, flux unfolded cross section is model dependent.

Reconstruction bias due to QE assumption is corrected under "RFG" model assumption.

One should be careful when comparing flux-unfolded data from different experiments.



5. CCQE double differential cross section

Flux-averaged double differential cross section (T_μ-cosθ)

This is the most complete information about neutrino cross section based on muon kinematic measurement.

The error shown here is shape error, a total normalization error (δN_T=10.8%) is separated.



5. CCQE double differential cross section

Flux-averaged double differential cross section (T_μ-cosθ)

This is the most complete information about neutrino cross section based on muon kinematic measurement.

The error shown here is shape error, a total normalization error (δN_T=10.8%) is separated.



6. Conclusions

Using the high statistics and high purity MiniBooNE ν_μ CCQE data sample (146,070 events, 26.5% efficiency, and 75.8% purity), the absolute cross section is measured. We especially emphasize the measurement of flux-averaged double differential cross section, because this is the most complete set of information for muon kinematics based neutrino interaction measurement. The double differential cross section is the model independent result.

We measured 35% higher cross section than RFG model with the world averaged nuclear parameter. Interesting to note, our total cross section is consistent with RFG model with nuclear parameters extracted from shape-only fit in our Q^2 data.

BooNE collaboration

University of Alabama
Bucknell University
University of Cincinnati
University of Colorado
Columbia University
Embry Riddle Aeronautical University
Fermi National Accelerator Laboratory
Indiana University
University of Florida
Los Alamos National Laboratory
Louisiana State University
Massachusetts Institute of Technology
University of Michigan
Princeton University
Saint Mary's University of Minnesota
Virginia Polytechnic Institute
Yale University

Moltes Grácies!

(¡Muchas Gracias!)

Back up

1. CCQE event measurement in MiniBooNE

cut type	efficiency
1. veto hits < 6 for all subevents	45.1
2. 1st subevent time T is in beam window	44.7
3. 1st subevent reconstructed vertex < 500 cm	37.5
4. 1st subevent kinetic energy > 200MeV	32.7
5. μ to e log likelihood cut	31.3
6. 2 subevent total	29.0
7. μ-e vertex distance cut	26.5

26.5% cut efficiency
75.8% purity
146,070 events with
5.58E20POT

4. CCQE absolute cross section

Absolute flux-averaged differential cross section formula

i :true index
j : reconstructed index

The cross section is function of true value, for example, $d\sigma^2/T_\mu/\cos\theta_\mu$, $d\sigma/dQ^2_{QE}$, etc

Integrated flux is removed, so it is called flux-averaged cross section

$$\sigma_i = \frac{\sum_j U_{ij}(d_j - b_j)}{\varepsilon_i (\Phi T)}$$

U_{ij} :unsmearing matrix

d_j :data vector

b_j :predicted background

T :integrated target number

ε_i :efficiency

Φ :integrated ν-flux

4. CCQE absolute cross section

Absolute flux-unfolded total cross section formula

i :true index
j : reconstructed index

The cross section is function of true neutrino energy, $\sigma[E_\nu^{QE}]$

Flux shape is removed bin by bin, so it is called flux-unfolded cross section

$$\sigma_i = \frac{\sum_j U_{ij}(d_j - b_j)}{\varepsilon_i (\Phi T)}$$

U_{ij} :unsmearing matrix

d_j :data vector

b_j :predicted background

T :integrated target number

ε_i :efficiency

Φ_i :ν-flux vector

5. CCQE flux error

Flux error

The flux error dominates total normalization error.

The shape error is weak, except high energy region, where HARP measurement has large error and skin effect of horn has large error.



5. CCQE background cross section error

Background cross section error

The background cross section error is small, because of high purity and in situ background constraint.

The large error comes from pion absorption, so the kinematic space of CC1π events has large error



5. CCQE detector error

Detector error

The detector error has the largest contribution to the shape error because it is related with the energy scale of muon.

However the contribution to the total normalization error is not so large.



Jon Link, Nov. 18, 2005
Fermilab Wine & Cheese seminar

E 23, NUMBER 11 1 JUNE 1981

Quasielastic neutrino scattering: A measurement of the weak nucleon axial-vector form factor

N. J. Baker, A. M. Cnops,* P. L. Connolly, S. A. Kahn, H. G. Kirk, M. J. Murtagh, R. B. Palmer, N. P. Samios, and M. Tanaka

Brookhaven National Laboratory, Upton, New York 11973

(Received 12 February 1981)

Brookhaven AGS 7ft D_2 Bubble Chamber

The quasielastic reaction $\nu_\mu n \to \mu^- p$ was studied in an experiment using the BNL 7-foot deuterium bubble chamber exposed to the wide-band neutrino beam with an average energy of 1.6 GeV. A total of 1138 quasielastic events in the momentum-transfer range $Q^2 = 0.06 - 3.00$ $(\mathrm{GeV}/c)^2$ were selected by kinematic fitting and particle identification and were used to extract the axial-vector form factor $F_A(Q^2)$ from the Q^2 distribution. In the framework of the conventional $V-A$ theory, we find that the dipole parametrization is favored over the monopole. The value of the axial-vector mass M_A in the dipole parametrization is 1.07 ± 0.06 GeV, which is in good agreement with both recent neutrino and electroproduction experiments. In addition, the standard assumptions of conserved vector current and no second-class currents are checked.

We have used a maximum likelihood method to extract M_A from the shape of the Q^2 distribution for each observed neutrino energy. This likelihood function $\mathcal{L}^I$ is independent of the shape of the neutrino spectrum . . .

They didn't even try to determine their ν flux from pion production and beam dynamics.

Phys. Rev. D 25, 617 (1982)

The distribution of events in neutrino energy for the 3C $\nu d \to \mu^- p p_s$ events is shown in Fig. 4 together with the quasielastic cross section $\sigma(\nu n \to \mu^- p)$ calculated using the standard $V-A$ theory with $M_A = 1.05 \pm 0.05$ GeV and $M_V = 0.84$ GeV. The absolute cross sections for the CC interactions have been measured using the quasielastic events and its known cross section.[4]

In subsequent cross section analyses the theoretical ("known") quas-ielastic cross section and observed quasi-elastic events were used to determine the flux.

Jon Link, Nov. 18, 2005
Fermilab Wine & Cheese seminar

REVIEW LETTERS 12 JULY 1982

Fermilab
15ft D_2 Bubble Chamber

Neutrino Flux and Total Charged-Current Cross Sections in High-Energy Neutrino-Deuterium Interactions

T. Kitagaki, S. Tanaka, H. Yuta, K. Abe, K. Hasegawa, A. Yamaguchi, K. Tamai, T. Hayashino, Y. Ohtani, and H. Hayano
Tohoku University, Sendai 980, Japan

To obtain the total cross section from the number of events, the neutrino flux has to be measured on an absolute scale. In this analysis, we determine the neutrino flux using 362 quasielastic events identified in our data[10] and the cross section for reaction (2) derived from the $V-A$ theory.

Again, they use QE events and theoretical cross section to calculate the ν.

When they try to get the flux from meson (π and K) production and decay kinematics they fail miserably for E_ν<30 GeV.



FIG. 2. Neutrino flux distribution obtained from the quasielastic events and the predicted cross section with M_A = 1.05 GeV. The solid curve is obtained from the best fit to the flux data for E_ν > 30 GeV. The dashed curve is taken from the Monte Carlo simulation of the flux.

Jon Link, Nov. 18, 2005
Fermilab Wine & Cheese seminar

ME 34, NUMBER 1 1 JULY 1986

Determination of the neutrino fluxes in the Brookhaven wide-band beams

L. A. Ahrens, S. H. Aronson, P. L. Connolly,* B. G. Gibbard, M. J. Murtagh, S. J. Murtagh,†
S. Terada, and D. H. White
Physics Department, Brookhaven National Laboratory, Upton, New York 11973

Brookhaven
AGS
Liquid Scintillator

The beam calculations described here were based on the Grote, Hagedorn, and Ranft (GHR) (Ref. 11) parametrization; that of Sanford and Wang was used for comparison. An estimate was made of pion production by reinteracting protons guided by the shape of the observed ν_μ spectrum and the observed angular distribution of muons from quasielastic events. The procedure is described[12] in the Appendix.

The Procedure

- Pion production cross sections in some low momentum bins are scaled up by 18 to 79%.
- The K^+ to π^+ ratio is increased by 25%.
- Overall neutrino (anti-neutrino) flux is increased by 10% (30%).

All driven by the neutrino events observed in the detector!

Jon Link, Nov. 18, 2005
Fermilab Wine & Cheese seminar

16, NUMBER 11 1 DECEMBER 1977

Argonne ZGS
12ft D_2 Bubble Chamber

Study of neutrino interactions in hydrogen and deuterium: Description of the experiment and study of the reaction $\nu + d \rightarrow \mu^- + p + p_s$ †

S. J. Barish,* J. Campbell,‡ G. Charlton,§ Y. Cho, M. Derrick, R. Engelmann,‖ L. G. Hyman, D. Jankowski, A. Mann,¶ B. Musgrave, P. Schreiner, P. F. Schultz, R. Singer, M. Szczekowski,** T. Wangler, and H. Yuta††
Argonne National Laboratory, Argonne, Illinois 60439

Flux derived from pion production data. Were able to test assumptions about the form of the cross section using absolute rate and shape information.

TABLE IV. Results of axial-form-factor fits.

Likelihood function	M_A^{Dipole} (GeV)	M_A^{Monopole} (GeV)	M_A^{Tripole} (GeV)
Rate	$0.75^{+0.13}_{-0.11}$	$0.45^{+0.11}_{-0.07}$	$0.96^{+0.17}_{-0.14}$
Shape	1.010 ± 0.09	0.56 ± 0.08	1.32 ± 0.11
Rate and shape	0.95 ± 0.09	0.52 ± 0.08	1.25 ± 0.11
Flux independent	0.95 ± 0.09	0.53 ± 0.08	1.25 ± 0.11

- Pion production measured in ZGS beams were used in this analysis
- A very careful job was done to normalize the beam.
- Yet they have a 25% inconsistency between the axial mass they measure considering only rate information verses considering only spectral information.

Interpretation: Their normalization is wrong.