


# First Anti-Neutrino oscillation results from MiniBooNE

Žarko Pavlović  
Los Alamos National Laboratory

# LSND signal

- MiniBooNE designed to test LSND
- LSND observed excess of  $\bar{\nu}_e$  in  $\bar{\nu}_\mu$  beam  
 $87.9 \pm 22.4 \pm 6.0$  ( $3.8\sigma$ )
- Assuming two neutrino oscillation:

$$P(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) = \sin^2(2\theta) \sin^2\left(\frac{1.27 L \Delta m^2}{E}\right)$$
$$= 0.245 \pm 0.067 \pm 0.045 \%$$



# LSND anomaly

- Within Standard Model 3 neutrino framework LSND signal is not compatible with atmospheric and solar neutrino data



$\nu_3$

$$\Delta m_{23}^2 = m_2^2 - m_3^2$$




$\nu_2$

$$\Delta m_{12}^2 = m_1^2 - m_2^2$$




$\nu_1$

$$\Delta m_{13}^2 = \Delta m_{12}^2 + \Delta m_{23}^2$$




# MiniBooNE



- Similar L/E as LSND
- MB detector:
  - 800 ton mineral oil Čerenkov detector
  - 12m diameter
  - 1280 inner PMT's, and 240 outer veto PMT's
- Focus either positive or negative mesons
- Collected & analysed:
  - $6.5e20$  POT in neutrino mode
  - $3.4e20$  POT in anti-neutrino mode



# Neutrino events

- Čerenkov rings used to identify the products of  $\nu$  interactions
- Looking for  $\nu_e$  CCQE signal events
- Backgrounds:
  - Intrinsic  $\nu_e$
  - Mis-identified events
- Use  $\nu_\mu$  CCQE to constrain intrinsic  $\nu_e$



# Neutrino run results

- No excess of events in signal region
- Ruled out  $2 \nu$  oscillations as source of LSND signal (assuming no CP or CPT violation)




A. A. Aguilar-Arevalo et al., Phys.Rev.Lett. 98: 231801,2007;  
A. A. Aguilar-Arevalo et al., Phys.Rev.Lett.102:101802,2009

# Neutrino run results

- Excess of events observed at low energy:  
 $128.8 \pm 20.4 \pm 38.3$  ( $3.0\sigma$ )
- Shape not consistent with 2  $\nu$  oscillations
- Magnitude consistent with LSND

- Anomaly Mediated Neutrino-Photon Interactions at Finite Baryon Density: Jeffrey A. Harvey, Christopher T. Hill, & Richard J. Hill, arXiv:0708.1281
- CP-Violation 3+2 Model: Maltoni & Schwetz, arXiv:0705.0107; T. Goldman, G. J. Stephenson Jr., B. H. J. McKellar, Phys. Rev. D75 (2007) 091301.
- Extra Dimensions 3+1 Model: Pas, Pakvasa, & Weiler, Phys. Rev. D72 (2005) 095017
- Lorentz Violation: Katori, Kostelecky, & Tayloe, Phys. Rev. D74 (2006) 105009
- CPT Violation 3+1 Model: Barger, Marfatia, & Whisnant, Phys. Lett. B576 (2003) 303
- New Gauge Boson with Sterile Neutrinos: Ann E. Nelson & Jonathan Walsh, arXiv: 0711.1363




# Anti-Neutrino run

- Further insight in low energy excess
- Provides direct test of LSND
- Statistics low at present
  - First result with  $3.4E20$  POT
  - The total number of CCQEs down by order of magnitude compared to neutrino run
- By the end of run this summer will have  $5E20$  POT
  - Approved for additional  $5E20$  POT run
- Same analysis as for neutrino mode with few checks


# Antineutrino results

- No significant excess observed at either low or high energy



# Limit from anti-neutrino run

- No significant excess observed at either low or high energy
- Assumes that neutrinos don't oscillate, only antineutrinos do
- Both LSND and null signal consistent with data at 90% CL
- Need more data



# Low Energy Excess

- Preliminary results of the fits to  $\nu$  and  $\bar{\nu}$  data in low E region
- Ignoring what we know about various backgrounds and looking only at how compatible two datasets are
- Preferred model: excess is due to neutrinos in the beam (no contribution from antineutrinos)

	<b>Stat. Error Only</b>	<b>100% Correlated Syst. Error</b>	<b>Uncorrelated Syst. Error</b>
Same $\nu, \bar{\nu}$ NC	0.1%	0.1%	6.7%
NC $\pi^0$ scaled	3.6%	6.4%	21.5%
POT scaled	0.0%	0.0%	1.8%
Bkgd scaled	2.7%	4.7%	19.2%
CC scaled	2.9%	5.2%	19.9%
Low-E Kaons	0.1%	0.1%	5.9%
$\nu$ scaled	38.4%	51.4%	58.0%

Maximum  $\chi^2$  probability from fits to  $\nu$  and  $\bar{\nu}$  data in 200-475 MeV range

# Conclusion

- MiniBooNE observes excess of low energy events in neutrino run consistent in magnitude with LSND signal, however energy shape not consistent with 2  $\nu$  oscillations
- With  $3.4e20$  POT MiniBooNE does not observe low energy excess in anti neutrino data
- High energy data consistent with both LSND best fit point and no oscillations
- Collecting more anti-neutrino data (approved for a new  $5e20$ POT run)