Recent results of and future plans for the Booster Neutrino Beamline

Kendall Mahn, Columbia University
for the MiniBooNE and SciBooNE collaborations

Outline

- One beam...
  - Booster Neutrino Beamline
  - Resulting flux
- Two experiments...
  - SciBooNE detectors
  - MiniBooNE detector
- Physics!
  - Oscillation
  - Cross section measurements
    - Neutrino and Antineutrino
- Summary

The big picture



- 8.9 GeV/c protons from Booster accelerator
- protons hit a target within a magnetic focusing horn and produce mesons
- The mesons decay into neutrinos the ~450 m decay region
- Neutrinos are observed in MiniBooNE and SciBooNE

Booster Neutrino Beamline (BNB)

- Booster
  - 4×10^{12} protons / 1.6 μs pulse delivered at up to 5 Hz
- Target
  - 1.7 λ Be target
- Magnetic horn
  - Pulses at 2.5 kV, 174kA
  - Increases flux by ~x6







Secondary meson production

HARP 8.9 GeV/c pBe π^+ production



- For π^+, K^+ ,and K^0 production use a parameterization to fit the existing data
- Errors span spread in data as well as fit errors

External measurements of pBe K^+ production from 9.5 to 24 GeV, scaled to 8.9 GeV/c



Neutrino flux

- Geant 4 model of beamline
- ν_μ predominantly from π^+ except at high energy (K^+)
- 0.5 % ν_e content
  - 52% $\mu \Rightarrow e^+ \nu_\mu \nu_e$
  - 29% $K^+ \Rightarrow \pi^+ e^+ \nu_e$
  - 14% $K^0 \Rightarrow \pi e \nu_e$
  - 5% other
- 6% antineutrino content



Antineutrino flux





- Reversible horn current; can focus negatively charged mesons for an antineutrino beam

  $\pi^- \Rightarrow \bar{\nu}_\mu$ vs $\pi^+ \Rightarrow \nu_\mu$
- Flux for antineutrino mode has substantial neutrino contamination, lower overall rate

Interactions in the BNB

- Nuance Monte Carlo
  D. Casper, NPS, 112 (2002) 161
- Largest interaction is charged current quasi elastic (CCQE)
  Common channel for normalization, oscillation analyses
- Next largest is charged current single pion production (CC π^+)
  background in CCQE samples
- Also Neutral current pion production (NC π^0) and Neutral current elastic scattering (NC EL)



Outline

- One beam...
  - Booster Neutrino Beamline
  - Resulting flux
- Two experiments...
  - SciBooNE detectors
  - MiniBooNE detector
- Physics!
  - Oscillation
  - Cross section measurements
    - Neutrino and Antineutrino
- Summary





SciBooNE

- Preexisting (free!) fine grained tracking detectors
- Insert into a running, modeled beamline
- Cross section measurements
  - Similar energy as T2K
  - Near detector for MiniBooNE
  - Measurement of rare and antineutrino cross sections
- Intent to run 1e20 pot in neutrino and antineutrino mode



The SciBooNE Collaboration

T. Ishii, M. Tanaka
High Energy Accelerator Research Organization (KEK)

T. Katori, C. Polly, R. Tayloe
Indiana University

Y. Hayato
Institute for Cosmic Ray Research (ICRR)

I.J. Taylor, Y. Uchida, J. Walding, D. Wark, M.O. Wascko(*)
Imperial College London

S.-I. Gomi, K. Hiraide, H. Kawamuko, H. Kubo, Y. Kurimoto, Y. Nakajima, T. Nakaya (*), K. Matsuoka, M. Taguchi, H. Tanaka, M. Yokoyama
Kyoto University

J. Alcaraz, G. Jover, F. Sanchez
Universitat Autonoma de Barcelona

R. Johnson
University of Cincinnati

M. Wilking, E.D. Zimmerman
University of Colorado, Boulder

A.A. Aguilar-Arevalo, L. Bugel, Z. Djurcic, J.M. Conrad, K.B.M. Mahn, V. Nguyen, M.H. Shaevitz, G.P. Zeller
Columbia University

R.H. Bernstein, S.J. Brice, B.C. Brown, D.A. Finley, T. Kobilarcik, A.D. Russell, R. Stefanski, R.J. Tesarek, H. White
Fermi National Accelerator Laboratory

W.C. Louis, R. Van de Water
Los Alamos National Laboratory

W. Metcalf
Louisiana State University

R. Napora
Purdue University Calumet

U. Dore, C. Giganti, P.F. Loverre, L. Ludovici, C. Mariani
Universita degli Studi di Roma "La Sapienza"

P. Nienaber
Saint Mary's University of Minnesota

Y. Miyachi,T.-A. Shibata, H. Takei
Tokyo Institute of Technology

J. Catala, A. Cervera-Villanueva, J.J. Gomez-Cadenas, M. Sorel, A. Tornero
Unversidad de Valencia

SciBooNE: SciBar



- 15 ton extruded scintillator (carbon target) with wavelength shifting fiber (WLS) readout
- Each cell is 2.5 x 1.3 x 300 cm
- 224 64 channel Multi-Anode PMT (14336 channels)
- Built for K2K



SciBooNE: Electron Catcher (EC)

- "spaghetti" calorimeter
  - 1mm scintillating fibers in grooves between lead foil
  - 256 channels
- 11 X_0 long
- 1 vertical and 1 horizontal plane; PMT readout at both ends
- Built for CHORUS, used for K2K

SciBooNE: Muon Range Detector (MRD)

- 12 5cm iron plates interspersed with scintillator counters; ~48 tons of material
- 13 horizontal and 12 vertical planes with 26 or 30 scintillator counters attached, each with a single PMT readout (362 channels)
- Assembled from scratch and spare parts at FNAL

Events in SciBooNE



- Tracking detector
  - Can use dE/dX to distinguish pion from proton from muon
  - proton tracks >10 cm are reconstructable
  - electrons stop in EC
  - Muons < 0.9 GeV stop in MRD
- Resolution
  - SciBar: 0.08 GeV muon energy, 1 degree angular resolution
  - EC: 14% $\sqrt{E}$
  - MRD: ~10% $\sqrt{E}$

(Real) Events in SciBooNE



CCQE $\overline{\nu}_\mu$



CC π^+ ν_μ

Timeline





- Mar. 2005: K2K ends
- Summer 2005: collaboration formed
- Nov. 2005: Proposed
- Dec. 2005: Approved, detectors sent from KEK
- July 2006: SciBar and EC arrive, MRD construction begins

Timeline

- Sept. 2006: Groundbreaking for new detector hall
- Nov. 2006 - Feb 2007: Assembly and construction of all 3 detectors



Timeline

- March 2007: Testing of detectors with cosmic rays
- April 2007: Detectors lowered into new detector hall

Timeline



Fermilab

Fermi National Accelerator Laboratory
P.O. Box 500 • Batavia, Il. • 60510-0500
630-840-3211 FAX 630-840-2900

Director's Office

July 20, 2007

Prof. Tsuyoshi Nakaya
High Energy Physics Group
Dept of Physics, Faculty of Science
Kyoto University
Kitashirakawa Oiwake-cho
Sakyo-ku, Kyoto, 606-8502
Japan

Dr. Morgan Wascko
Physics Department
Imperial College London
Blackett Lab 525
Prince Consort Road
London SW7 2AZ
United Kingdom

Dear Tsuyoshi and Morgan,

Given your progress on the SciBooNE experiment, including a formal Memorandum of Understanding between SciBooNE and the Laboratory, I am pleased to grant formal Stage II approval to the experiment.

I wish you successful continuation of the data taking.

Sincerely,

Piermaria Oddone



- May 31st, 2007: First data events in all 3 detectors!
- July 20th, 2007: Received Stage II approval

Outline

- One beam...
  - Booster Neutrino Beamline
  - Resulting flux
- Two experiments...
  - SciBooNE detectors
  - MiniBooNE detector
- Physics!
  - Oscillation
  - Cross section measurements
    - Neutrino and Antineutrino
- Summary



The MiniBooNE Collaboration

Y.Liu, D.Perevalov, I.Stancu
University of Alabama
S.Koutsoliotas
Bucknell University
R.A.Johnson, J.L.Raaf
University of Cincinnati
T.Hart, R.H.Nelson, M.Tzanov
M.Wilking, E.D.Zimmerman
University of Colorado
A.A.Aguilar-Arevalo, L.Bugel L.Coney,
J.M.Conrad, Z. Djurcic,
K.B.M.Mahn, J.Monroe, D.Schmitz
M.H.Shaevitz, M.Sorel, G.P.Zeller
Columbia University
D.Smith
Embry Riddle Aeronautical University
L.Bartoszek, C.Bhat, S.J.Brice
B.C.Brown, D. A. Finley, R.Ford, F.G.Garcia,
P.Kasper, T.Kobilarcik, I.Kourbanis, A.Malensek,
W.Marsh, P.Martin, F.Mills, C.Moore, E.Prebys,
A.D.Russell , P.Spentzouris,
R.J.Stefanski, T.Williams
Fermi National Accelerator Laboratory
D.C.Cox, T.Katori, H.Meyer, C.C.Polly
R.Tayloe
Indiana University
G.T.Garvey, A.Green, C.Green, W.C.Louis, G.McGregor, S.McKenney
G.B.Mills, H.Ray, V.Sandberg, B.Sapp, R.Schirato, R.Van de Water
N.L.Walbridge, D.H.White
Los Alamos National Laboratory
R.Imlay, W.Metcalf, S.Ouedraogo, M.O.Wascko
Louisiana State University
J.Cao, Y.Liu, B.P.Roe, H.J.Yang
University of Michigan
A.O.Bazarko, P.D.Meyers, R.B.Patterson, F.C.Shoemaker, H.A.Tanaka
Princeton University
P.Nienaber ***Saint Mary's University of Minnesota***
J. M. Link ***Virginia Polytechnic Institute***
E.Hawker ***Western Illinois University***
A.Curioni, B.T.Fleming ***Yale University***

MiniBooNE: Detector

800 ton mineral oil (carbon target) Cherenkov detector

- 12 m diameter sphere
- 1280 inner region 8” PMTs, 10% photocathode coverage
- 240 outer ‘veto’ region PMTs

Resolution:

- Charge: 1.4 PE
- Time: 1.7 ns



Events in MiniBooNE

Use hit topology, timing to determine event type

- Outgoing lepton implies flavor of neutrino for charged current events
- Reconstructed quantities: track length, angle relative to beam direction
- Fundamental: timing, charge of hits, early/late hit fractions
- Geometry: position from wall of tank



Additional information in scintillation light

- ~25% of the light in the tank due to mineral oil
- Unlike prompt Cherenkov light, scintillation light is delayed
- Amount depends on particle type





Subevents in MiniBooNE



19.2 μsec trigger window around 1.6μsec beam

Trigger on neutrino event ($\nu_\mu + n \Rightarrow \mu + p$) initially, subsequent electron from muon decay ($\mu \Rightarrow e^+ \nu_\mu \nu_e$)

Each cluster of hits in time is a 'subevent'

Precuts in MiniBooNE



Apply precuts to eliminate 10 kHz cosmic ray background



Minimal veto activity (veto hits < 6)

First subevent with tank hits above decay electron endpoint (tank hits > 200)



Calibrating MiniBooNE



Model of light propagation in mineral oil

Dominant light source is well understood Cherenkov light

Also must model:

- Scintillation
  - yield, spectrum, decay times
- Fluorescence (absorption and reemision of Cherenkov light)
  - rate, spectrum, decay times
- Scattering
  - Rayleigh, Raman, Particulate (Mie)
- Absorption
- Reflection
  - tank walls, PMT faces
- PMT effects
  - single pe charge response, charge linearity



External measurements

- Scintillation from p beam (IUCF)
- Scintillation from cosmic μ (Cincinnati)
- Fluorescence Spectroscopy (FNAL)
- Time resolved spectroscopy (JHU, Princeton)
- Attenuation (Cincinnati)

Internal measurements

- Cosmic muons and decay electrons, Laser flasks

Outline

- One beam...
  - Booster Neutrino Beamline
  - Resulting flux
- Two experiments...
  - SciBooNE detectors
  - MiniBooNE detector
- Physics!
  - Oscillation
  - Cross section measurements
    - Neutrino and Antineutrino
- Summary



MiniBooNE

- Purpose: confirm or refute LSND
  - LSND observed an excess of $\bar{\nu}_e$ in a $\bar{\nu}_\mu$ beam; assuming 2 parameter mixing gives a 0.25% oscillation probability
  - Run with same L/E, different L, E, signal signature
- Started in 1998, oscillation results this year
  - 5.57e20 protons on target in neutrino mode
  - 2.2e20 protons on target and counting in antineutrino mode



LSND Beam Excess

Oscillation Analysis



ν_μ

0.5% intrinsic ν_e

ν_e selection cuts



E_ν(QE)

Signal

(Δm^2=1.2 eV^2, $\sin^2 2\theta$=0.003)

Background

- misidentified ν_μ (mainly π^0s)
- ν_e from μ^+ decay
- ν_e from K^+, K^0 decay
- $\Delta \Rightarrow N\gamma$
- Out of tank events ('dirt')

Do the ν_μ oscillate into ν_e ?

- Produce ν_μ
- Select ν_e
- Observe an excess or not?

ν_e selection cuts: particle identification (PID)

Two PID algorithms used

- Likelihood based analysis (TBL): e/μ, e/π^0 and $m_{\pi 0}$ cuts
- A "boosted decision tree" (BDT) algorithm to separate e, μ, π^0



"Likelihood" method:

- Each event is a combination of hits, charge, and position
- Form Q, T pdfs, fit for 7 track parameters to distinguish electron-muon hypothesis

ν_e selection cuts: particle identification (PID)

Two PID algorithms used

- Likelihood based analysis (TBL): e/μ, e/π^0 and $m_{\pi 0}$ cuts
- A "boosted decision tree" (BDT) algorithm to separate e, μ, π^0

A decision tree is similar to a neural net

- Cut first on the variable which gives the most separation of signal to background, at the point where it gives the most separation. Then cut on next best variable...



"Boosting" is a method to additionally separate signal from background, by weighting events

- Increase weight of misclassified events in current tree, and remake tree. Repeat ~100-1000x. Sum all the trees, by counting events on signal leaves as +1, and -1 otherwise. This forms the PID variable.

Oscillation Analysis



ν_μ

0.5% intrinsic ν_e

ν_e selection cuts



Signal E_ν(QE)

(Δm^2=1.2 eV^2, $\sin^2 2\theta$=0.003)

Background

- misidentified ν_μ (mainly π^0s)
- ν_e from μ^+ decay
- ν_e from K^+, K^0 decay
- $\Delta \Rightarrow N\gamma$
- Out of tank events ('dirt')

Do the ν_μ oscillate into ν_e ?

- Produce ν_μ
- Select ν_e
- Observe an excess or not?

NC π^0 tagging

- Asymmetric decays where only one photon is observed look just like a single electron, or a CCQE ν_e event
- Select 1 subevent, minimal veto activity, above decay endpoint, and within fiducial volume
- Create two likelihood variables-- 1 ring vs 2 ring hypothesis, and 1 ring electron or muon like
- Select events which fit well to 2 ring, electron like, and which fall within the reconstructed π^0 mass peak
- very pure (~90%) sample









NC π^0 rate measurement

- Compare the observed π^0 rate to the MC as a function of π^0 momentum, and make a correction factor
- Reweight the misidentified π^0s based on their momentum by this correction factor
- This is also the correction applied to the $\Delta \Rightarrow N\gamma$ events for the oscillation analysis

$M_{\gamma\gamma}$ Mass Distribution for Various $p_{\pi 0}$ Momentum Bins



Error 'budget'

source of uncertainty on nue background	TBL/BDT % error	constrained by MB data?	Reduced by relating ν_μ to ν_e
Flux from π^+/μ^+ decay	6.2 / 4.3	Y	Y
Flux from K^+ decay	3.3 /1.0	Y	Y
Flux from K^0 decay	1.5 / 0.4	Y	Y
Target/Beam model	2.8 / 1.3	Y	Y
ν cross section	12.3 / 10.5	Y	Y
NC π^0 yield	1.8 / 1.5	Y	
Out of tank events	0.8 / 3.4	Y	
Optical Model	6.1 / 10.5	Y	Y
DAQ electronics model	7.5 / 10.8	Y	

* shows errors before ν_e / ν_μ constraint is applied

All of our errors are highly correlated, but here are the diagonal errors

Constrain all samples with MiniBooNE data

Link between ν_e and ν_μ samples further reduces errors, like a 'near to far ratio'

Result



- TBL analysis shows no excess in analysis region, but excess at low energy
- Excess cannot be described based on LSND and a simple 2 ν mixing hypothesis; still under investigation

- BDT also has no sign of excess in analysis region
- Also shows an excess at low energy, but errors are larger and data low relative to prediction elsewhere complicates the issue



Sensitivity



- TBL limit (solid), BDT (dash)
- Agreement between independent analyses
- Incompatible with LSND at 98% CL at all Δm^2 for 2 neutrino mixing hypothesis

Outline

- One beam...
  - Booster Neutrino Beamline
  - Resulting flux
- Two experiments...
  - MiniBooNE detector
  - SciBooNE detectors
- Physics!
  - Oscillation
  - Cross section measurements
    - Neutrino and Antineutrino
- Summary

Cross sections around 1 GeV

World's data on neutrino cross sections vs Eν



- Nothing measured to better than 15-20%
- Many measurements not on nuclear targets
- Can take ratios of processes to CCQE to avoid some flux uncertainties
- Or measure same process in two detectors

Cross sections around 1 GeV

World's data on antineutrino cross sections vs Eν



BNB, T2K
mean Eν (GeV)

- Need antineutrino cross sections for future experiments
  - Compare oscillation in neutrino and antineutrino modes to map out CP violation and mass hierarchy in neutrino mixing



P(osc) vs δCP, for normal and inverted hierarchies

Charged Current Quasi-Elastic (CCQE) interaction



$$Q^2 = -m_\mu{}^2 + 2E_\nu(E_\mu - p_\mu \cos\theta_\mu)$$

- Tag single muon events and their decay electron
  - 2 subevents, minimal veto activity in both
  - muon-like track, 2nd event below decay electron energy endpoint
  - both events within fiducial volume
  - 74% purity, 197k event sample
- Simple relationship between neutrino, and lepton measurables makes this a ‘golden’ mode for oscillation measurements

CCQE cross section formalism

C. H. Llewellyn Smith, Phys. Rep. 3C, 261 (1972)

$$< N'|J_\mu|N > \; = \; \overline{u}(N')\left[\gamma_\mu \underline{F'_V(q^2)} + \frac{i\sigma_{\mu\nu}q^\nu \xi \underline{F^2_V(q^2)}}{2M} + \gamma_5\gamma_\mu \underline{F_A(q^2)}\right] u(N)$$

- QE cross section can be written in as a function of Q^2, in terms of vector (F_V) and axial vector (F_A) form factors
- F_V are constrained by electron scattering experiments; F_A by neutrino scattering only
- Assuming a dipole form:

$$F_A(Q^2) = \frac{g_A}{(1 + Q^2 / M_A{}^2)^2}$$

where M_A is the 'axial mass'

- This picture is complicated by the presence of nuclear effects, strong at low Q^2

CCQE measurement



T. Katori

Simultaneous shape fit for M_A and nuclear effect parameter (κ) in Q^2 with a relativistic Fermi Gas model

M_A= 1.23 +/- 0.20 GeV,
κ = 1.019 +/- 0.011

(hep-ex/0706.0926)

Recent K2K SciBar result:

M_A= 1.14 +/- 0.11 GeV

World Average: M_A= 1.03 +/- 0.02 GeV (J.Phys. G28, R1 (2002)

NC π^0:Coherent fraction

- π^0 can be produced either from the excitation of a delta (resonant), or from the excitation of the whole nucleus (coherent)
- Coherent production is forward peaked, and a larger component of antineutrino running



J. Link

resonant



coherent



Will convert rate measurement into a cross section
Make a measurement of the coherent fraction:
Coh/Res = (19.5 +/-1.1 (stat) +/- 2.5 (sys))%

Antineutrinos: Coherent fraction

- π^0 can be produced either from the excitation of a delta (resonant), or from the excitation of the whole nucleus (coherent)
- Coherent production is forward peaked, and a larger component of antineutrino running

coherent included



No coherent component



V. Nguyen

resonant



coherent



- Preliminary antineutrino sample fits better to a nonzero coherent fraction

NC Elastic



Signature is scintillation light of proton below Cherenkov threshold

- 1 subevent, minimal veto activity
- Cutting on late light fraction eliminates the Cherenkov light from low energy electron events
- Radial cut to reduce events from surrounding dirt which don't fire veto
- ~84% purity

Only 1 other measurement made so far (overlaid here)

Sensitive to the same M_A as CCQE



D. Cox

Prospects: SciBooNE and MiniBooNE

- SciBooNE has superior efficiency and reconstruction capabilities, however, MiniBooNE dominates with statistics
- Leverage each other
  - ν_μ disappearance: measure initial rate of events in SciBooNE, compare directly to MiniBooNE
- MiniBooNE: tags CC π^+ with 3 subevents (1 muon/pion, and two decay electrons)
- SciBooNE: reconstruct entire final state!
  - If pion is absorbed, will affect kinematics





Prospects: SciBooNE and MiniBooNE

- SciBooNE has superior efficiency and reconstruction capabilities, however, MiniBooNE dominates with statistics
- Antineutrino running:
  - MiniBooNE can select CC π^+ (neutrino only process) in antineutrino beam to measure neutrino contamination
  - SciBooNE can tag on an event by event basis



CCQE ν_μ



CCQE $\overline{\nu_\mu}$

Summary

Lots of physics done with the Booster Neutrino Beam

- Recent MiniBooNE oscillation result
- Cross section results from MiniBooNE (CCQE)

And more to come!

- Soon NC π^0 and NC elastic, and CC π^+
- ν_μ disappearance

SciBooNE proposed, installed, and running

- Will complement MiniBooNE's capabilities
- Map out cross sections around 1 GeV in both neutrino, and antineutrino interactions

Backup slides

Building an error matrix: light propagation in detector

Create different 'universes' with the parameters varied within errors

Compare them to muon decay electron (Michel) sample variables, such as time, charge, hit topology

Keep universes which have a good $\chi 2$ as compared to data

Michel electron t distribution



This restricted space defines the parameters and correlations. Draw from the new space, and build an error matrix:

$$M_{ij}^{OM} = \frac{1}{universes} \sum_{k=1}^{universes} (N_{cv} - N_k)_i (N_{cv} - N_k)_j$$



Out of tank ("dirt") events

- Events from interactions in the surrounding rock produce photons which pass the veto and give events within the inner tank (so called "dirt") events
- Create a sample of enhanced dirt events
  - in time with beam, minimal veto activity, 1 subevent, not decay electron
  - low energy, high radius
- Checks prediction spatial distribution, energy spectrum of these events; sets the normalization for dirt events in the ν_e sample



visible energy (MeV)

Constraint Example: ν_e from μ^+

Without employing a link between ν_e and ν_μ , ν_e from μ^+ would have all aforementioned errors: flux, cross section, detector uncertainties

However, for each ν_e produced from a μ^+, there was a corresponding ν_μ and we observe that ν_μ spectrum

This is true here because the pion decay is very forward

Therefore, we know that some combination of cross sections, flux, etc errors are excluded by our own data, and so the error is reduced





25 m absorber problem

Two absorber plates, suspended above the decay volume, fell

First drop after March shutdown, second in August 2006, 10% rate drop for each

Chains were rated at 17 tons each, 2 chains supported each 25 ton plate

Clean break point on chain (not weld point); metal had become brittle

25 METER ABSORBER - ELEVATION & CROSS-SECTION